



King's Research Portal

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Patton, R., & Marshall, E. (2014). Physician heal thyself: Results from a single hospital pilot survey of alcohol consumption among general hospital patients and staff. *Journal of studies on alcohol and drugs*, 75(1), 189-190. http://www.jsad.com/jsad/downloadarticle/Physician_Heal_Thyself_Results_From_a_SingleHospital_Pilot_Survey_of_AI/5245.pdf

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

CORRESPONDENCE

“Physician Heal Thyself”: Results From a Single-Hospital Pilot Survey of Alcohol Consumption Among General Hospital Patients and Staff

Dear Editor:

Excessive alcohol consumption and its associated harms remain as a major public health issue in England (NHS Information Centre, 2011). Despite a plethora of evidence demonstrating that screening for misuse, followed up by some intervention (information booklet, simple advice, or counseling), can significantly reduce both consumption and consequences (Kaner et al., 2013), the implementation of alcohol identification and brief advice (IBA) across Health Service settings has been limited (Drummond et al., 2011). One possible reason for this might be the complex relationship between our own alcohol consumption and attitudes toward offering help and advice to others (Huntley et al., 2004).

To what extent might current levels of consumption among hospital staff account for clinical inertia regarding alcohol IBA? On a single day, ward managers at a large Inner London teaching hospital distributed our alcohol use survey to all staff, patients, and visitors. The survey contained items on respondent demographics and smoking status as well as the Alcohol Use Disorders Identification Test–consumption (AUDIT-C) alcohol questions. For patients, a single item queried their perception of the role that alcohol may have played in necessitating their presence at the hospital.

A total of 360 completed questionnaires were returned (72% female). Responses were received from patients (54%), staff (40%), and visitors (6%). Age ranged from 16 to 90 years, with an average of 43. Patient responses came from inpatient (45%), outpatient (45%), and the emergency department (10%) areas.

The majority of respondents consumed alcohol on one or more occasions per month (72.5%). A total of 133 (37%) scored 5 or more on the AUDIT-C and were identified as hazardous drinkers (33% of all males, 39% of all females), and, of these, 20 (5.6% of sample, 15% of hazardous drinkers) met criteria for harmful/dependent drinking (score of ≥ 8 , based on thresholds established by Rubinsky et al., 2010). As might be expected, patients based in the emergency department had the highest proportion of hazardous drinkers (44%) compared with those in the inpatient (36%)

and outpatient (26%) settings. A significantly higher proportion of hospital staff were identified as hazardous drinkers compared with patients, 44.4% vs. 31.4%, $\chi^2(2) = 6,046$, $p < .05$; however, there were no significant differences between staff and patients on harmful/dependent drinking status (4.2% vs. 5.8%).

We do not have reliable data on the number of questionnaires distributed, and therefore we are unable to calculate the response rate of our survey or estimate how representative our sample was of the wider population. Nevertheless, our finding that a significantly higher proportion of staff than patients was hazardous drinkers is of interest. We suggest that, given the now-established correlation between personal alcohol consumption and clinical inertia (Geirsson, 2013), particular attention should be paid to the provision of help and advice about drinking for those already working within health services, as this may well have a positive effect on the wider implementation of patient-focused alcohol IBA programs and should also help reduce levels of hazardous and harmful consumption among health service employees themselves.

ROBERT PATTON, PH.D.
Addictions Department
King's College London
London, Southwark, United Kingdom
robert.patton@kcl.ac.uk

JANE MARSHALL, M.B., B.CH.
Addictions Department
King's College London
London, Southwark, United Kingdom
Jane.Marshall@slam.nhs.uk

References

- Drummond, C., Gual, A., Goos, C., Godfrey, C., Deluca, P., Von Der Goltz, C., . . . Kaner, E. (2011). Identifying the gap between need and intervention for alcohol use disorders in Europe. *Addiction*, 106, Supplement 1, 31–36.

- Geirsson, M. (2013). *Alcohol prevention in Swedish primary health care: Staff knowledge about risk drinking and attitudes towards working with brief alcohol intervention. Where do we go from here?* Gothenburg, Sweden: University of Gothenburg. Retrieved from https://gupea.ub.gu.se/bitstream/2077/26274/1/gupea_2077_26274_1.pdf
- Huntley, J. S., Patton, R., & Touquet, R. (2004). Attitudes towards alcohol of emergency department doctors trained in the detection of alcohol misuse. *Annals of the Royal College of Surgeons of England*, 86, 329–333.
- Kaner, E., Bland, M., Cassidy, P., Coulton, S., Dale, V., Deluca, P., . . . Drummond, C. (2013). Effectiveness of screening and brief alcohol intervention in primary care (SIPS trial): Pragmatic cluster randomised controlled trial. *BMJ*, 346, e8501. Retrieved from <http://www.bmj.com/content/346/bmj.e8501>
- NHS Information Centre. (2011). *Statistics on Alcohol: England, 2011*. London, England: The Health and Social Care Information Centre. Retrieved from <https://catalogue.ic.nhs.uk/publications/public-health/alcohol/alco-eng-2011/alco-eng-2011-rep.pdf>
- Rubinsky, A. D., Kivlahan, D. R., Volk, R. J., Maynard, C., & Bradley, K. A. (2010). Estimating risk of alcohol dependence using alcohol screening scores. *Drug and Alcohol Dependence*, 108, 29–36.